Carbohydrate Research, 229 (1992) C19–C21
Elsevier Science Publishers B.V., Amsterdam

CONTENTS

Publisher's Note v

The crystal structure of the inclusion complex of cyclomaltoheptaose (β-cyclodextrin) with 4-*tert*-butyltoluene
I.M. Mavridis and E. Hadjoudis (Athens, Greece) 1

The crystal and molecular structures of the hexitol hexa-acetates
J. Kopf, M. Morf, B. Zimmer (Hamburg, Germany), M. Bischoff and P. Köll (Oldenburg, Germany) 17

The crystal structure of α-D-glucopyranosyl potassium hydrogenphosphate
T. Lis (Wrocław, Poland) 33

Molecular and crystal structure of konjac glucomannan in the mannan II polymorphic form
T. Yui, K. Ogawa and A. Sarko (Syracuse, NY, USA) 41

Molecular and crystal structure of the regenerated form of (1 → 3)-α-D-mannan
T. Yui, K. Ogawa and A. Sarko (Syracuse, NY, USA) 57

Synthesis and conformational analysis of substituted 2,6-dioxabicyclo[3.1.1]heptanes: 1,3-anhydro-2,4-di-*O*-benzyl-6-deoxy-β-D-glucopyranose by ^{1}H-NMR spectroscopy and molecular mechanics calculation
X. Wu, F. Kong, D. Lu and P. Zhang (Beijing, China) 75

Addition and cyclisation reactions of sugar phenylhydrazone derivatives via a 1,4-elimination process: preparation of 3-(2-acetamido-1-phenylhydrazonobutyl)-2(1*H*)-quinoxalinone and 3-(1-acetamidopropyl)- pyrazolo[3,4-*b*]quinoxaline (flavazole) derivatives
L. Somogyi (Debrecen, Hungary) 89

Stereoselective syntheses of a di-, tri-, and tetra-saccharide fragment of *Shigella dysenteriae* type 1 O-antigen using 3,4,6-tri-*O*-acetyl-2-azido-2-deoxy-α-D-glucopyranosyl chloride as a glycosyl donor
V. Pavliak, P. Kováč and C.P.J. Glaudemans (Bethesda, MD, USA) 103

Synthesis of 2-*O*-(2-acetamido-2-deoxy-β-D-glucopyranosyl)-5a-carba-α-D-mannopyranose, and 5a-carba-2-*O*- and 3,6-di-*O*-(α-D-mannopyranosyl)-α-D-mannopyranoses
S. Ogawa and K. Nishi (Yokohama, Japan) 117

Synthesis and characterization of highly substituted deoxyfluorocellulose acetate
N. Kasuya, K. Iiyama and A. Ishizu (Tokyo, Japan) 131

Transacetalation: a convenient, nonaqueous method for effecting the deprotection of isopropylidene and benzylidene derivatives of sugars
M.A. Andrews and G.L. Gould (Upton, NY, USA) 141

Notes

Phase transfer catalyzed synthesis of 4-nitrophenyl 1-thio-β-D-glycobiosides
F.D. Tropper, F.O. Andersson, C. Grand-Maître and R. Roy (Ottawa, ON, Canada) 149

Synthesis of a repeating pentasaccharide fragment of the capsular polysaccharide of *Streptococcus pneumoniae* type 18C
A.M.P. Van Steijn, J.G.M. Van der Ven, P. Van Seeventer, J.P. Kamerling and J.F.G. Vliegenthart (Utrecht, Netherlands) 155

Synthesis of methyl 2-*C*-acetamidomethyl-2-deoxy-α-D-glucopyranoside and its *manno* isomer
P. Sarda, A. Olesker and G. Lukacs (Gif-sur-Yvette, France) 161

Synthesis of a chiral polysubstituted *cis*-fused decalin containing an angular *C*-methyl group by an intramolecular Diels–Alder reaction
S. Rémy, F. Jaekel, A. Olesker and G. Lukacs (Gif-sur-Yvette, France) 167

Preparation of building blocks for carba-oligosaccharides related to cell-surface glycans
S. Ogawa, T. Tonegawa, K. Nishi and J. Yokoyama (Yokohama, Japan) 173

Synthesis of feruloylated and *p*-coumaroylated methyl glycosides
R.F. Helm, J. Ralph and R.D. Hatfield (Madison, WI, USA) 183

Preliminary communications

Synthesis of 1-deoxynojirimycin-containing glycans related to the Lewis X and sialyl-Lewis X epitopes recognized by LEC-CAMs
H. Furui, M. Kiso and A. Hasegawa (Gifu, Japan) C1

Methyl-3-*O*-(2-acetamido-2-deoxy-6-thio-β-D-glucopyranosyl)-β-D-galactopyranoside: a slow reacting acceptor-analogue which inhibits glycosylation by UDP-D-galactose–*N*-acetyl-D-glucosamine-(1→4)-β-D-galactosyltransferase
Y. Kajihara, H. Hashimoto and H. Kodama (Yokohama, Japan) C5

Papers

A conformational study of α-L-Rha*p*-(1 → 2)-α-L-Rha*p*-(1 → OMe) by NMR nuclear Overhauser effect spectroscopy (NOESY) and molecular dynamics calculations
G. Widmalm, R.A. Byrd and W. Egan (Bethesda, MD, USA) 195

A nuclear magnetic resonance spectroscopic investigation of Kdo-containing oligosaccharides related to the genus-specific epitope of *Chlamydia* lipopolysaccharides
K. Bock, J.U. Thomsen (Valby, Denmark), P. Kosma, R. Christian (Vienna, Austria), O. Holst and H. Brade (Borstel, Germany) 213

Karplus-type equation for vicinal carbon–proton coupling constants for the C–S–C–H pathway in 1-thioglycosides
I. Tvaroska (Bratislava, Czechoslovakia), K. Mazeau, M. Blanc-Muesser, S. Lavaitte, H. Driguez and F.R. Taravel (Grenoble, France) 225

The use of neocarrabiose oligosaccharides with different length and sulphate substitution as model compounds for ^{1}H-NMR spectroscopy
S.H. Knutsen and H. Grasdalen (Trondheim, Norway) 233

NMR spectroscopy and conformational analysis of substituted 1,2:5,6-di-*O*-isopropylidene-α-D-allofuranose derivatives
D.C. Lankin, S.T. Nugent and S.N. Rao (Skokie, IL, USA) 245

Binding studies on internal immunodeterminants: synthesis of β-(1 → 6)-linked oligosaccharide methyl glycosides having one to four internal D-galactopyranosyl residues flanked by gentiobiose residues
T. Ziegler (Stuttgart, Germany), H. Sutoris and C.P.J. Glaudemans (Bethesda, MD, USA) 271

Synthesis of derivatives of methyl 3-deoxy-3-*C*-formyl- α-D-*arabino*-pentofuranoside
S. Benefice-Malouet, P.L. Coe and R.T. Walker (Birmingham, United Kingdom) 293

Influence of malondialdehyde on the Maillard degradation of Amadori compounds
A. Gómez-Sánchez, I. Hermosín and I. Maya (Seville, Spain) 307

Synthesis of D-glucose 3- and 6-[2-(perfluoroalkyl)ethyl phosphates]: a new type of anionic surfactant for biomedical use
A. Milius, J. Greiner and J.G. Riess (Nice, France) 323

Interaction of linear manno-oligosaccharides with three mannose-specific bulb lectins. Comparison with mannose/glucose-binding lectins
H. Kaku and I.J. Goldstein (Ann Arbor, MI, USA) 337

New mannose-specific lectins from garlic (*Allium sativum*) and ramsons (*Allium ursinum*) bulbs
H. Kaku, I.J. Goldstein (Ann Arbor, MI), E.J.M. Van Damme and W.J. Peumans (Leuven, Belgium) 347

Anticoagulant and antithrombin activities of oversulfated fucans
T. Nishino and T. Nagumo (Kanagawa, Japan) 355

Notes

Reaction of 2-deoxy-6-*O*-[2,3-dideoxy-4,6-*O*-isopropylidene-2,3-(*N*-tosylepimino)-α-D-mannopyranosyl]- 4,5-*O*-isopropylidene-1,3-di-*N*-tosylstreptamine with potassium hydrogenfluoride
Y. Kobayashi, N. Taneichi, T. Tsuchiya and K. Koga (Kawasaki, Japan) 363

Epimerization of reducing terminal groups of (1 → 2)-linked D-gluco- and D-manno-disaccharides in aqueous sodium hydroxide
H. Kobayashi, N. Shibata, S. Konno, K. Hisamichi and S. Suzuki (Miyagi, Japan) 369

Announcement C11

Corrigendum C12

Author index C13

Subject index C15

Contents (vol. 229) C19